

ON TIGHT PROJECTIVE DESIGNS

YU. I. LYUBICH

ABSTRACT. It is shown that among all tight designs in $\mathbb{F}\mathbb{P}^n \neq \mathbb{R}\mathbb{P}^1$, where $\mathbb{F}$ is $\mathbb{R}$ or $\mathbb{C}$, or $\mathbb{H}$ (quaternions), only 5-designs in $\mathbb{C}\mathbb{P}^1$ [14] have irrational angle set. This is the only case of equal ranks of the first and the last irreducible idempotent in the corresponding Bose-Mesner algebra.

Keywords: projective design, angle set, Bose-Mesner algebra.

AMS Classification: 05B30

1. INTRODUCTION

A well known theorem of Bannai and Hoggar [3] states that there are no tight t -designs in $\mathbb{F}\mathbb{P}^n \neq \mathbb{R}\mathbb{P}^1$ if $t \geq 6$. Moreover, a theorem of Hoggar [10] states the same for $t \geq 4$ if $\mathbb{F} \neq \mathbb{R}$. Surprisingly, a tight 5-design in $\mathbb{C}\mathbb{P}^1$ has been constructed in [14], so Hoggar's theorem has to be corrected. The results of [3] and [10] are essentially based on Theorem 2.6(c) [9] that states that the angle set of every tight t -design in $\mathbb{F}\mathbb{P}^n \neq \mathbb{R}\mathbb{P}^1$ is rational. But it is not rational for the 5-design constructed in [14].

In the present paper we investigate this contradiction and prove that *the only cases where the angle set is not rational are*

- (1) $\mathbb{F} = \mathbb{C}$, $n = 1$, $t = 5$ and
- (2) $\mathbb{F} = \mathbb{R}$, $n = 1$, $t \neq 1, 2, 3, 5$.

A fortiori, there are no complications in [3] where $t \geq 6$ by assumption.

Our principal observation is that if $t = 2s - 1$, $s \geq 2$ then the last irreducible idempotent L_s in the corresponding Bose-Mesner algebra is not E_s from the proof of Theorem 2.6(c) [9] (actually, from [18]). Nevertheless, $\text{rk } L_s \neq \text{rk } E_1$, except for our case (1). This "critical inequality" implies the rationality of the angle set, similarly to the argument in [9]. This material is concentrated in Section 4 of the present paper, while Sections 2 and 3 contain all the necessary background and preliminary analysis.

2. PROJECTIVE t -DESIGNS

For the reader's convenience we basically use the same notation as in [8] and other related papers. Let us recall this notation. In particular, let

$$\mathbb{F} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}; \quad m = \frac{1}{2}(\mathbb{F} : \mathbb{R}) = \begin{cases} 1/2 & \mathbb{F} = \mathbb{R} \\ 1 & \mathbb{F} = \mathbb{C} \\ 2 & \mathbb{F} = \mathbb{H} \end{cases}; \quad N = m(n+1).$$

The number $2N$ is nothing but the real (topological) dimension of the $\mathbb{F}$ -linear space $\mathbb{F}^{n+1}$. The latter consists of all $(n+1) \times 1$ matrices (columns) over $\mathbb{F}$ with the standard addition and multiplication by scalars $\tau \in \mathbb{F}$ from the right (for

definiteness). As usual, the inner product of $a, b \in \mathbb{F}^{n+1}$ is a^*b where a^* is the row conjugate transpose to a . Accordingly, the set

$$S^{2N-1} = \{a : a^*a = 1\}$$

is the unit sphere in $\mathbb{F}^{n+1}$. A quotient set of the sphere with respect to the equivalence relation $a_1 \sim a_2 \iff a_1 = a_2\lambda$, $\lambda \in \mathbb{F}$, $|\lambda| = 1$, is the projective space $\mathbb{FP}^n$. The “inner product” $(\hat{a}, \hat{b}) = |a^*b|^2$ in $\mathbb{FP}^n$ is well-defined through the natural mapping $a \mapsto \hat{a}$ from S^{2N-1} onto $\mathbb{FP}^n$. Obviously, $(\hat{b}, \hat{a}) = (\hat{a}, \hat{b})$ and $0 \leq (\hat{a}, \hat{b}) \leq 1$ with the equality $(\hat{a}, \hat{b}) = 1$ if and only if $\hat{a} = \hat{b}$. For every nonempty $X \subset \mathbb{FP}^n$ its angle set is

$$A(X) = \{(x, y) : x, y \in X, x \neq y\}$$

The related combinatorial parameters are

$$s = |A(X)|, \quad e = |A(X) \setminus \{0\}|, \quad \epsilon = s - e = |A(X) \cap \{0\}|.$$

Let $P_i^{(\alpha, \beta)}(\tau)$ be the Jacobi polynomials [20] such that

$$\deg P_i^{(\alpha, \beta)} = i, \quad P_i^{(\alpha, \beta)}(1) = \frac{(\alpha + 1)_i}{i!} \quad (2.1)$$

where

$$(\alpha + 1)_i = \prod_{l=1}^i (\alpha + l), \quad (\alpha + 1)_0 = 1.$$

In particular, $P_0^{(\alpha, \beta)}(\tau) \equiv 1$. In what follows we fix

$$\alpha = N - m - 1, \quad \beta = m - 1 \quad (2.2)$$

and set

$$P_i(\xi) = P_i^{(\alpha, \beta)}(2\xi - 1), \quad (2.3)$$

for short. A finite nonempty subset $X \subset \mathbb{FP}^n$ is called a t -design if

$$\sum_{x \in X} P_i((x, y)) = 0, \quad y \in X, \quad 1 \leq i \leq t. \quad (2.4)$$

Let X be a t -design and let

$$R_e^\epsilon(\xi) = \frac{(N)_s}{(m)_s} P_e^{(\alpha+1, \beta+\epsilon)}(2\xi - 1).$$

In particular,

$$R_e^\epsilon(1) = \frac{(N)_s (N - m + 1)_e}{(m)_s e!} \quad (2.5)$$

The following theorems are fundamental, see [1], [2], [11]. (Cf. [6] for the spherical designs.)

Theorem A. *The inequalities*

$$t \leq s + e, \quad |X| \geq R_e^\epsilon(1)$$

hold, and the equalities

$$t = s + e, \quad |X| = R_e^\epsilon(1)$$

are equivalent.

In the latter case the t -design X is called *tight*. Note that $t = s + e$ is equivalent to $e = \lfloor t/2 \rfloor$, $\epsilon = \text{res}_2(t)$.

Theorem B. *If X is a tight t -design then $A(X)$ coincides with the set of roots of the polynomial $\xi^\epsilon R_e^\epsilon(\xi)$.*

Recall that these roots are simple and lie on $(0, 1)$.

Theorem C. *Let X be a subset of $\mathbb{FP}^n$ such that $|X| = R_e^\epsilon(1)$ and $A(X)$ coincides with the set of roots of $\xi^\epsilon R_e^\epsilon(\xi)$, then X is a tight $(2e + \epsilon)$ -design.*

The projective t -designs can be characterized as the averaging sets in the sense of [19] for suitable spaces of functions on $\mathbb{FP}^n$. Usually, these spaces are described in terms of harmonic analysis but we prefer a more elementary approach [15], [16].

We say that a mapping $\phi : \mathbb{FP}^n \rightarrow \mathbb{C}$ is a *polynomial function* if it is of the form

$$\phi(\hat{a}) = \psi(a), \quad a \in S^{2N-1},$$

where ψ is a polynomial on $\mathbb{F}^{n+1}$ in real coordinates. This ψ must be invariant with respect to the rotations of $\mathbb{F}$, i.e. $\psi(a\lambda) = \psi(a)$ for all $\lambda \in \mathbb{F}$, $|\lambda| = 1$. It is not unique but becomes unique if it is required to be homogeneous (which is always possible) of minimal degree. The latter is said to be the *degree* of ϕ . The number $\deg \phi$ is an even integer since $\psi(-a) = \psi(a)$.

Example 2.1. For every $t \in \mathbb{N}$ and every $y \in \mathbb{FP}^n$ the function $\phi_{2t;y}(x) = (x, y)^t$, $x \in \mathbb{FP}^n$, is a polynomial function of degree $2t$.

Given $d \in 2\mathbb{N}$, we denote by $\text{Pol}_{\mathbb{F}}(d)$ the space of all polynomial functions of degrees $\leq d$. It has been proven in [16] that the family $\{\phi_{d;y} : y \in \mathbb{FP}^n\}$ spans the whole space $\text{Pol}_{\mathbb{F}}(d)$. We apply this result to prove the following

Proposition 2.2. *A finite nonempty set $X \subset \mathbb{FP}^n$ is a tight t -design if and only if*

$$\frac{1}{|X|} \sum_{x \in X} \phi(x) = \int_{S^{2N-1}} \tilde{\phi}(a) d\sigma(a), \quad \phi \in \text{Pol}_{\mathbb{F}}(2t), \quad (2.6)$$

where $\tilde{\phi}$ is induced by the natural mapping $S^{2N-1} \rightarrow \mathbb{FP}^n$ and σ is the normalized Lebesgue measure.

Proof. The identity (2.6) is equivalent to

$$\frac{1}{|X|} \sum_{x \in X} F((x, y)) = \int_{S^{2N-1}} F(|a^* b|^2) d\sigma(a), \quad (2.7)$$

where $y = \hat{b}$, $b \in S^{2N-1}$, F runs over the space Π_t of all univariate polynomials of degrees $\leq t$. By a known integration formula (see [8], Theorem 2.11) one can rewrite (2.7) in the form

$$\frac{1}{|X|} \sum_{x \in X} F((x, y)) = \int_{-1}^1 F\left(\frac{1+\tau}{2}\right) \Omega_{\alpha,\beta}(\tau) d\tau, \quad \psi \in \Pi_t, \quad (2.8)$$

where $\Omega_{\alpha,\beta}(\tau)$ is the normalized Jacobi weight, i.e.

$$\Omega_{\alpha,\beta}(\tau) = c_{\alpha,\beta} (1-\tau)^\alpha (1+\tau)^\beta, \quad -1 < \tau < 1, \quad (2.9)$$

with

$$c_{\alpha,\beta} = \left(\int_{-1}^1 (1-\tau)^\alpha (1+\tau)^\beta d\tau \right)^{-1} = \frac{\Gamma(\alpha + \beta + 2)}{2^{\alpha+\beta+1} \Gamma(\alpha + 1) \Gamma(\beta + 1)}. \quad (2.10)$$

In turn, (2.8) is equivalent to its restriction to $F = P_i(\xi)$, $1 \leq i \leq t$, since these polynomials constitute a basis in Π_t . It remains to note that

$$\int_{-1}^1 P_i \left(\frac{1+\tau}{2} \right) \Omega_{\alpha,\beta}(\tau) d\tau = \int_{-1}^1 P_i^{(\alpha,\beta)}(\tau) \Omega_{\alpha,\beta}(\tau) d\tau = 0$$

by (2.3) and the $\Omega_{\alpha,\beta}$ -orthogonality of the system $\{P_i^{(\alpha,\beta)} : i \geq 0\}$. $\square$

Corollary 2.3. *Let $X \subset \mathbb{FP}^n$ be a t -design. Then*

$$\frac{1}{|X|} \sum_{x \in X} P((u, x)) Q((x, v)) = \int_{S^{2N-1}} P(|a^* c|^2) Q(|c^* b|^2) d\sigma(c) \quad (2.11)$$

for $u = \hat{a}$, $v = \hat{b}$ and all univariate polynomials P, Q such that $\deg P + \deg Q \leq t$.

Proof. The mapping $x \mapsto P((u, x)) Q((x, v))$, $x \in \mathbb{FP}^n$, is a polynomial function of degree $\leq 2t$. $\square$

Corollary 2.4. *Let X, P, Q be fixed under the conditions of Corollary 2.3. Then the value*

$$\sum_{x \in X} P((u, x)) Q((x, v))$$

depends only on the inner product (u, v) of $u, v \in \mathbb{FP}^n$.

Proof. Let $(u_1, v_1) = (u, v)$, i.e. $|a_1^* b_1|^2 = |a^* b|^2$ where $\hat{a}_1 = u_1$, $\hat{b}_1 = v_1$. Without loss of generality one can assume that $a_1^* b_1 = a^* b$. Then there exists a $(n+1) \times (n+1)$ matrix T over $\mathbb{F}$ such that $T^* T = \text{id}$ and $a_1 = Ta$, $b_1 = Tb$. This substitution in (2.11) is equivalent to the change of variable $c \mapsto T^* c$. The latter does not affect the integral since the measure σ is orthogonally invariant. $\square$

3. BOSE-MESNER ALGEBRA

Let X be a finite nonempty subset of $\mathbb{FP}^n$ and let

$$A'(X) = A(X) \cup \{1\} = \{(x, y) : x, y \in X\},$$

so that $|A'(X)| = s + 1$. The $X \times X$ matrices of the form

$$M_F = [F((x, y))]_{x, y \in X} \quad (3.1)$$

where F runs over all functions $A'(X) \rightarrow \mathbb{C}$, constitute a complex linear space $\mathcal{D}(X)$. Its natural basis consists of the matrices

$$\Delta_\zeta = [\delta_{\zeta, (x, y)}]_{x, y \in X}, \quad \zeta \in A'(X), \quad (3.2)$$

thus, $\dim \mathcal{D}(X) = s + 1$. The Lagrange interpolation formula allows us to let F in (3.1) run over the polynomial space Π_s , so that we have the isomorphism $F \mapsto M_F$ between Π_s and $\mathcal{D}(X)$. In particular, if $F|_{A(X)} = 0$ and $F(1) = 1$ then $M_F = I$, the unit matrix.

According to Corollary 2.4 for $P, Q \in \Pi_s$, the matrix product $M_P M_Q$ belongs to $\mathcal{D}(X)$ if $\deg P + \deg Q \leq t$. However, this condition is not fulfilled if $t = 2s - 1$ and $\deg P = \deg Q = s$. Moreover, *Corollary 2.4 cannot be extended to this situation if X is tight*. Indeed, suppose to the contrary that

$$\sum_{x \in X} (u, x)^s (x, v)^s = \Phi((u, v)) \quad (u, v \in \mathbb{FP}^n)$$

with a function $\Phi : [0, 1] \rightarrow \mathbb{R}_+$. Setting $v = u$ we obtain

$$\sum_{x \in X} (u, x)^{2s} = \Phi(1), \quad u \in \mathbb{F}\mathbb{P}^n.$$

In other words,

$$\sum_{c \in \tilde{X}} |a^* c|^{4s} = \Phi(1), \quad a \in S^{2N-1},$$

where $\tilde{X} \subset S^{2N-1}$ is a complete system of representatives of points $x \in X$, $|\tilde{X}| = |X|$. By integration over a we obtain

$$\Phi(1) = \left(\int_{S^{2N-1}} |a^* c|^{4s} d\sigma(a) \right) \cdot |X|$$

since the integral does not depend on c . As a result,

$$\frac{1}{|X|} \sum_{x \in X} \phi_{4s;u}(x) = \int_{S^{2N-1}} \tilde{\phi}_{4s;u}(a) d\sigma(a), \quad (3.3)$$

and by linearity, (3.3) extends to the whole space $\text{Pol}_{\mathbb{F}}(4s)$. Thus, X is a $2s$ -design which is a contradiction since $2s = t + 1$.

Nevertheless, under the constraint $u, v \in X$, one can extend Corollary 2.4 to $t = 2s - 1$ and P, Q such that $\max(\deg P, \deg Q) = s$. This follows from the construction of a basis in $\mathcal{D}(X)$ using the Jacobi polynomials (cf. [6], Remark 7.6).

Lemma 3.1. *Let X be a t -design in $\mathbb{F}\mathbb{P}^n$ and let $s = \lfloor \frac{t+1}{2} \rfloor$. Then $s + 1$ matrices $M_i = M_{P_i}$, $0 \leq i \leq s$, constitute a basis $\mathcal{M}$ of $\mathcal{D}(X)$ such that*

$$M_i M_k = |X| M_i \delta_{ik} \rho_{\mu(i,k)} \quad (3.4)$$

where $\mu(i, k) = \min(i, k)$ and all $\rho_j > 0$.

Proof. The matrices M_i are linearly independent because of the linear independence of the polynomials P_i . Since $|\mathcal{M}| = s + 1$, this is a basis of $\mathcal{D}(X)$. Now note that

$$\int_{S^{2N-1}} P_i(|a^* c|^2) P_k(|c^* b|^2) d\sigma(c) = 0, \quad i \neq k,$$

by the addition formula for polynomial functions [15] (cf. [7], [13], [17]). The same formula with $i = k$ yields

$$\int_{S^{2N-1}} P_i(|a^* c|^2) P_i(|c^* b|^2) d\sigma(c) = \chi_i P_i(|a^* b|^2) \quad (3.5)$$

where $\chi_i > 0$. Assuming $\mu(i, k) \leq s - 1$ (a fortiori, $i + k \leq 2s - 1 \leq t$) and using Corollary 2.3 we get (3.4) with

$$\rho_j = \chi_j, \quad 0 \leq j \leq s - 1. \quad (3.6)$$

In particular, $M_i M_s = M_s M_i = 0$ for $0 \leq i \leq s - 1$. It remains to consider the case $i = k = s$.

If t is even the $t = 2s$ and Corollary 2.3 is applicable to $i = k = s$, so $M_s^2 = |X| M_s \rho_s$ with

$$\rho_s = \chi_s. \quad (3.7)$$

Let t be odd, so $t = 2s - 1$. Then we decompose the unity matrix I for the basis $\mathcal{M}$,

$$I = \sum_{i=0}^s \lambda_i M_i, \quad (3.8)$$

and get $M_s = \lambda_s M_s^2$ multiplying (3.8) by M_s . This yields

$$\lambda_s = \frac{\text{tr } M_s}{\text{tr } M_s^2} = \frac{|X|P_s(1)}{\sum_{x,y} P_s^2((x,y))} > 0, \quad (3.9)$$

and then $M_s^2 = |X|M_s\rho_s$ with

$$\rho_s = (\lambda_s|X|)^{-1}. \quad (3.10)$$

□

Remark 3.2. The formulas (3.6) and (3.7) are joined in

$$\rho_i = \chi_i, \quad 0 \leq i \leq [t/2], \quad (3.11)$$

while (3.10) appears only for $t = 2s - 1$ in addition to (3.11).

Remark 3.3. The multiplication table (3.4) shows that under conditions of Lemma 3.1 $\mathcal{D}(X)$ is a commutative matrix algebra, the *Bose-Mesner algebra of X* [4], [5], [6].

In what follows the conditions of Lemma 3.1 are assumed to be fulfilled. By setting

$$L_i = \frac{M_i}{\rho_i|X|} = \frac{1}{\rho_i|X|} [P_i((x,y))]_{x,y \in X} \quad (3.12)$$

the basis $\mathcal{M}$ turns into $\mathcal{L} = \{L_i\}_0^s$ consisting of idempotents ($L_i^2 = L_i$) which are pairwise orthogonal ($L_i L_k = 0$ for $i \neq k$). It is important to calculate their ranks.

We have

$$\text{rk } L_i = \text{tr } L_i = \rho_i^{-1} P_i(1), \quad 0 \leq i \leq s,$$

hence,

$$\text{rk } L_i = \chi_i^{-1} P_i(1) = \frac{P_i^2(1)}{\int_{S^{2N-1}} P_i^2(|a^*c|^2) d\sigma(c)}, \quad 0 \leq i \leq [t/2]$$

by (3.11) and (3.5) for $a = b$. Finally,

$$\text{rk } L_i = \frac{\left(P_i^{(\alpha,\beta)}(1)\right)^2}{\int_{-1}^1 \left(P_i^{(\alpha,\beta)}(\tau)\right)^2 \Omega_{(\alpha,\beta)}(\tau) d\tau}, \quad 0 \leq i \leq [t/2]. \quad (3.13)$$

In particular, $\text{rk } L_0 = 1$. In addition to (3.13) we have to find $\text{rk } L_s$ in the case $t = 2s - 1$. Formula (3.10) is not effective to this end since λ_s is unknown. Indeed, in (3.9) we cannot proceed to the formally corresponding integral in the denominator. Instead of this, we return to the decomposition of unity and express $\text{rk } L_s$ through $\text{rk } L_i$, $0 \leq i \leq s - 1$. We have

$$I = \sum_{i=0}^s L_i,$$

whence,

$$\text{rk } L_s = \text{tr } L_s = |X| - \sum_{i=0}^{s-1} \text{tr } L_i = |X| - \sum_{i=0}^{s-1} \text{rk } L_i.$$

By substitution from (3.13) the last sum can be written as $c_{\alpha,\beta}^{-1} K_{s-1}^{(\alpha,\beta)}(1,1)$, where $K_{s-1}^{(\alpha,\beta)}(\cdot, \cdot)$ is the reproducing kernel of the Jacobi polynomials with respect to the

weight $(1 - \tau)^\alpha(1 + \tau)^\beta$, see [20], Section 4.5. According to (2.10) and formula (4.5.8) from [20] we obtain

$$\text{rk } L_s = |X| - \frac{\Gamma(s + \alpha + \beta + 1)\Gamma(s + \alpha + 1)\Gamma(\beta + 1)}{\Gamma(\alpha + \beta + 2)\Gamma(\alpha + 2)\Gamma(s)}$$

With our α, β defined by (2.2)

$$\text{rk } L_s = |X| - \frac{\Gamma(N + s - 1)\Gamma(N - m + s)\Gamma(m)}{\Gamma(N)\Gamma(N - m + 1)\Gamma(m + s - 1)\Gamma(s)} = |X| - \frac{(N)_{s-1}(N - m + 1)_{s-1}}{(m)_{s-1}(s - 1)!} \quad (3.14)$$

Lemma 3.4. *Let X be a tight t -design in $\mathbb{FP}^n$ with $t = 2s - 1$. Then*

$$\text{rk } L_s = \frac{(N)_{s-1}(N - m)_s}{(m)_s(s - 1)!} \quad (3.15)$$

Proof. In this case $e = s - 1$, $\epsilon = 1$, so (2.5) yields

$$|X| = R_{s-1}^1(1) = \frac{(N)_s(N - m + 1)_{s-1}}{(m)_s(s - 1)!}$$

□

The ranks of the other L_i (including L_s if $t = 2s$) can be explicitly calculated by (3.13), (2.10) and (2.1) combined with (4.33) of [20]. This results in

Lemma 3.5. *Let X be a t -design in $\mathbb{FP}^n$ with $s = \lceil \frac{t+1}{2} \rceil$. Then*

$$\text{rk } L_i = \frac{(N)_{i-1}(N - m)_i(N + 2i - 1)}{(m)_i i!}, \quad 0 \leq i \leq \lfloor t/2 \rfloor. \quad (3.16)$$

Remark 3.6. Formula (3.16) yields the true value $\text{rk } L_0 = 1$ by setting $(\gamma - 1)(\gamma)_{-1} = 1$ for all γ .

Corollary 3.7. *The inequality*

$$\text{rk } L_i > \text{rk } L_{i-1}, \quad 1 \leq i \leq \lfloor t/2 \rfloor, \quad (3.17)$$

holds, except for $X \subset \mathbb{FP}^1$. In the latter case

$$\text{rk } L_i = 2, \quad 1 \leq i \leq \lfloor t/2 \rfloor. \quad (3.18)$$

Now note that our idempotents L_i coincide with the matrices E_i from [9] for $0 \leq i \leq \lfloor t/2 \rfloor$ but $L_s \neq E_s$ if $t = 2s - 1$, $X \not\subset \mathbb{FP}^1$. Indeed, according to (2.5) from [9],

$$E_i((x, y)) = \frac{1}{|X|} [Q_i((x, y))]_{x, y \in X}, \quad 0 \leq i \leq s, \quad (3.19)$$

where $Q_i(\xi)$ is proportional to $P_i(\xi)$ and

$$Q_i(1) = \frac{(N)_{i-1}(N - m)_i(N + 2i - 1)}{(m)_i i!}, \quad i \geq 0. \quad (3.20)$$

Hence, E_i are proportional to L_i for all i , $0 \leq i \leq s$. Moreover, if $0 \leq i \leq \lfloor t/2 \rfloor$ then $\text{tr } E_i = Q_i(1) = \text{tr } L_i$ by (3.20) and (3.16). Hence, $E_i = L_i$ for $0 \leq i \leq \lfloor t/2 \rfloor$. However, if $t = 2s - 1$ (so $s = \lfloor t/2 \rfloor + 1$) and $X \not\subset \mathbb{FP}^1$ then $\text{tr } E_s = Q_s(1) > \text{tr } L_s$, see (3.15). In this case $\text{tr } E_s > \text{rk } E_s$, so E_s is not an idempotent. This is an obstacle to the full proof of Theorem 2.6 [9] of the rationality of $A(X)$. To overcome this difficulty, it suffices to change E_s for L_s (when $t = 2s - 1$, $s \geq 2$) but then the “critical inequality” $\text{rk } L_s \neq \text{rk } L_1$ is needed. However, the latter is not always true. We clarify this intricate situation in the next section.

4. THE CRITICAL INEQUALITY AND RATIONALITY THEOREM

We prove the following

Theorem 4.1. *With $t = 2s - 1$, $s \geq 2$, the inequality*

$$\operatorname{rk} L_s \neq \operatorname{rk} L_1 \quad (4.1)$$

holds for every tight t -design $X \subset \mathbb{F}\mathbb{P}^n$, except for a tight 5-design in $\mathbb{C}\mathbb{P}^1$.

Proof. From (3.15) it follows that

$$\operatorname{rk} L_s \geq \frac{(N)_1(N-m)_2}{(m)_2 \cdot 1!} = \frac{N(N-m)(N-m+1)}{m(m+1)}$$

since the right side of (3.15) increases with s . On the other hand, (3.16) yields

$$\operatorname{rk} L_1 = \frac{(N)_0(N-m)_1(N+1)}{(m)_1 \cdot 1!} = \frac{(N-m)(N+1)}{m} \quad (4.2)$$

Hence,

$$\operatorname{rk} L_s - \operatorname{rk} L_1 \geq \frac{(N-m)((N-m)^2 - (m^2 + m + 1))}{m(m+1)}$$

Since $N - m = mn$ we obtain $\operatorname{rk} L_s > \operatorname{rk} L_1$ for $n^2 > 1 + m^{-1} + m^{-2}$, i.e. for $n \geq 3$ if $\mathbb{F} = \mathbb{R}$ and for $n \geq 2$ if $\mathbb{F} = \mathbb{C}$ or $\mathbb{H}$. It remains to consider two cases.

- (1) $\mathbb{F} = \mathbb{R}$, $n = 2$. Then $m = 1/2$, $N = 3/2$, so $\operatorname{rk} L_1 = 5$ by (4.2), while $\operatorname{rk} L_s = 2s$ by (3.15).
- (2) $\mathbb{F}$ is arbitrary, $n = 1$. Then $N = 2m$, hence

$$\operatorname{rk} L_s - \operatorname{rk} L_1 = \frac{(2m)_{s-1}}{(s-1)!} - (2m+1) = \begin{cases} -1 & \mathbb{F} = \mathbb{R} \\ s-3 & \mathbb{F} = \mathbb{C} \\ \frac{1}{6}s(s+1)(s+2) - 5 & \mathbb{F} = \mathbb{H} \end{cases}$$

We see that $\operatorname{rk} L_s \neq \operatorname{rk} L_1$ with the only exception $\mathbb{F} = \mathbb{C}$, $s = 3$, so $t = 5$. $\square$

A tight 5-design in $\mathbb{C}\mathbb{P}^1$ should contain 12 points since its parameters are $m = 1$, $N = 2$, $s = 3$, $e = 2$, $\epsilon = 1$. Accordingly, (2.5) becomes

$$R_2^1(1) = \frac{(2)_3(2)_2}{(1)_3 \cdot 2!} = 12.$$

Such a design has been constructed in [14] as the projective image of an orbit of the binary icosahedral group that is a subgroup of $SU(2)$. Its representatives on the unit sphere $S^3 \subset \mathbb{C}^2$ are

$$a_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, a_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, a_k = \begin{cases} \mu \begin{pmatrix} \lambda \eta^{k-3} \\ 1 \end{pmatrix} & 3 \leq k \leq 7, \\ \mu \begin{pmatrix} \eta^{k-3} \\ -\lambda \end{pmatrix} & 8 \leq k \leq 12, \end{cases}$$

where

$$\eta = \exp \frac{2\pi i}{5}, \quad \lambda = 2 \cos \frac{2\pi}{5} = \frac{\sqrt{5}-1}{2}, \quad \mu = \frac{1}{2 \sin \frac{4\pi}{5}} = \sqrt{\frac{5+\sqrt{5}}{10}}.$$

We omit an elementary calculation of the inner products $(x_j, x_k) = |a_j^* a_k|^2$, only noting that

$$\lambda^2 + \lambda - 1 = 0, \quad (2 - \lambda)\mu^2 = 1, \quad |\eta^r - 1|^2 = 0, 2 - \lambda, 3 + \lambda,$$

the latter for $r \equiv 0, \pm 1, \pm 2 \pmod{5}$, accordingly. A calculation yields,

$$A(X) = \left\{ 0, \frac{5 - \sqrt{5}}{10}, \frac{5 + \sqrt{5}}{10} \right\}, \quad (4.3)$$

so $s = 3$, $e = 2$, $\epsilon = 1$. The polynomial

$$\xi R_2^1(\xi) = 4\xi P_2^{(1,1)}(2\xi - 1) = 6\xi(5\xi^2 - 5\xi + 1)$$

annihilates $A(X)$. By Theorem C of Section 2 our X is indeed a tight 5-design.

The angle set (4.3) is not rational. This occurs because of the equality $\text{rk } L_3 = \text{rk } L_1$. In fact, $\text{rk } L_0 = 1$, $\text{rk } L_1 = 3$, $\text{rk } L_2 = 5$, $\text{rk } L_3 = 3$, by (3.15) and (3.16).

In the following corrected form of Theorem 2.6 [9] the case $X \subset \mathbb{RP}^1$ is also included for completeness. In this case, (3.18) makes it possible for $A(X)$ to be not rational, though $\text{rk } L_s \neq \text{rk } L_1$ under the conditions of Theorem 4.1.

Theorem 4.2. *Let X be a tight t -design in $\mathbb{FP}^n$. Then the angle set $A(X)$ is rational, except for two cases: 1) $X \subset \mathbb{CP}^1$, $t = 5$; 2) $X \subset \mathbb{RP}^1$, $t \neq 1, 2, 3, 5$.*

Proof. For $X \not\subset \mathbb{RP}^1$ the proof is the same as in [9] but with L_s instead of E_s when $t = 2s - 1$, $s \geq 2$, and using our Theorem 4.1 in this case.

Now let $X \subset \mathbb{RP}^1$. Then it is the projective image of a regular $(2t + 2)$ -gon as easily follows from [12]. Therefore,

$$A(X) \setminus \{0\} = \left\{ \cos^2 \frac{k\pi}{t+1} \right\}_1^e = \left\{ \frac{1}{2} \left(1 + \cos \frac{2k\pi}{t+1} \right) \right\}_1^e$$

where $e = [t/2]$. Since $\cos m\theta$ is a polynomial of $\cos \theta$ with integer coefficients, the set $A(X)$ is rational if and only if the number $\rho = \cos \frac{2\pi}{t+1}$ is rational. Obviously, the latter is true if $t = 1, 2, 3, 5$. Conversely, let $\rho \in \mathbb{Q}$. Then the complex number $w = \exp(2\pi i/(t+1))$ satisfies the equation $w^2 - 2\rho w + 1 = 0$. On the other hand, this is a primitive root of 1 of degree $t+1$. It is known that irreducible (over $\mathbb{Q}$) equation for w is of degree $\varphi(t+1)$ where φ is the Euler function. Hence, $\varphi(t+1) \leq 2$. If $\varphi(t+1) = 1$ then $t = 1$. If $\varphi(t+1) = 2$ then $t \in \{2, 3, 5\}$ as easily follows from the classical formula

$$\varphi \left(\prod_{i=1}^r q_i^{\nu_i} \right) = \prod_{i=1}^r q_i^{\nu_i-1} (q_i - 1)$$

where $q_1, \dots, q_r$ are prime divisors of $t+1$. □

REFERENCES

- [1] E. Bannai. On extremal finite sets in the sphere and other metric spaces. In *Algebraic, extremal and metric combinatorics, 1986 (Montreal, PQ, 1986)*, volume 131 of *London Math. Soc. Lecture Note Ser.*, pages 13–38. Cambridge Univ. Press, Cambridge, 1988.
- [2] E. Bannai and S. G. Hoggar. On tight t -designs in compact symmetric spaces of rank one. *Proc. Japan Acad. Ser. A Math. Sci.*, 61(3):78–82, 1985.
- [3] E. Bannai and S. G. Hoggar. Tight t -designs and squarefree integers. *European J. Combin.*, 10(2):113–135, 1989.
- [4] R. C. Bose and D. M. Mesner. On linear associative algebras corresponding to association schemes of partially balanced designs. *Ann. Math. Statist.*, 30:21–38, 1959.

- [5] P. Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Res. Rep. Suppl.*, (10):vi+97, 1973.
- [6] P. Delsarte, J. M. Goethals, and J. J. Seidel. Spherical codes and designs. *Geometriae Dedicata*, 6(3):363–388, 1977.
- [7] S. G. Hoggar. Zonal functions and the symplectic group. Preprint, Mat. Inst. Aarhus Univ. 1–22, 1977.
- [8] S. G. Hoggar. t -designs in projective spaces. *European J. Combin.*, 3(3):233–254, 1982.
- [9] S. G. Hoggar. Tight t -designs and octonions. *Mitt. Math. Sem. Giessen*, (165):1–16, 1984.
- [10] S. G. Hoggar. Tight 4- and 5-designs in projective spaces. *Graphs Combin.*, 5(1):87–94, 1989.
- [11] S. G. Hoggar. t -designs in Delsarte spaces. In *Coding theory and design theory, Part II*, volume 21 of *IMA Vol. Math. Appl.*, pages 144–165. Springer, New York, 1990.
- [12] Y. Hong. On spherical t -designs in $\mathbf{R}^2$. *European J. Combin.*, 3(3):255–258, 1982.
- [13] T. Koornwinder. The addition formula for Jacobi polynomials and spherical harmonics. *SIAM J. Appl. Math.*, 25:236–246, 1973.
- [14] Yu. I. Lyubich and O. A. Shatalova. Euclidean subspaces of the complex spaces l_p^n constructed by orbits of finite subgroups of $SU(m)$. *Geom. Dedicata*, 86(1-3):169–178, 2001.
- [15] Yu. I. Lyubich and O. A. Shatalova. Isometric embeddings of finite-dimensional l_p -spaces over the quaternions. *Algebra i Analiz*, 16(1):15–32, 2004.
- [16] Yu. I. Lyubich and O. A. Shatalova. Polynomial functions on the classical projective spaces. *Studia Math.*, 170(1):77–87, 2005.
- [17] C. Müller. *Spherical harmonics*, volume 17 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1966.
- [18] A. Neumaier. *Combinatorial configuration in terms of distances*. Memorandum 81-09, Dept. of Math. Eindhoven Univ., 1981.
- [19] P. D. Seymour and T. Zaslavsky. Averaging sets: a generalization of mean values and spherical designs. *Adv. in Math.*, 52(3):213–240, 1984.
- [20] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., 1975.

Department of Mathematics
 Technion, Haifa 32000, Israel
 email: lyubich@tx.technion.ac.il